



VAKSIMUNE® POX

Is used to protect chicken against Fowl Pox.
The vaccine is safe to be used in any age of chickens including grower, layer or breeder.

(Fowl Pox Beaudette Strain, Live Vaccine, Liquid Form)

Composition : Beaudette strain at least 10^3 EID₅₀/ dose

High degree of safety.

Propagated in
SPF 

Does not exhibit any side effect.

Long lasting stability.


Fowl Pox

Fowl pox is a slow spreading viral infection of chickens characterized by proliferative lesions in the skin (cutaneous form) that progress to form thick scabs and by lesions in upper gastro-intestinal and respiratory tracts (diphtheritic form)




Administration of **VAKSIMUNE® POX**

1. Shake the vaccine thoroughly before use.
2. Expose the chicken wing web.
3. Soak the needle into vaccine.
4. Prick the wing web with the needle containing the vaccine.
5. One dose of vaccine is equal to 0.01 ml.



How 'Takes' Appear

The usual 'Take' consists of some swelling at the site of the puncture as early as the fourth day following vaccination. The swelling increases during the next five days until a scab is formed. Revaccinate birds that do not show 'Takes'.



Vaccination Schedule

Primary : 4-6 Weeks age

Booster : 14-16 Weeks age (Optional)

Withdrawal Period : None

Storage : Store in Refrigerator at 2°C - 8°C .

